

Dissimilarity in the evolution of solar EUV and solar radio emission (2800 MHz) during 1999–2002

R. P. Kane

Instituto Nacional de Pesquisas Espaciais, São Jose dos Campos, Brazil

Received 30 January 2003; revised 9 September 2003; accepted 17 September 2003; published 24 December 2003.

[1] For solar UV and EUV the F10 (2800 MHz solar radio emission), Lyman alpha, and Mg II are used as proxies. During 1999–2002, many solar indices showed two maxima, one near July 2000 and another near January 2002. For F10 and Lyman alpha, the second maximum was higher than the first maximum by $\sim 10\%$, but for Rz, the second maximum was lower than the first maximum by $\sim 4\%$. The observed EUV (26–34 nm from SEM-SOHO) had the second maximum higher than the first maximum, but only by $\sim 3.5\%$. The model values given in SOLAR2000 [Tobiska *et al.*, 2000] show patterns of 26–34 nm qualitatively similar to F10, with second maximum larger than the first maximum, but the magnitude is much larger, $\sim 20\%$. Thus there is a discrepancy of $\sim 15\%$ between observed EUV and SOLAR2000 model EUV during 2001–2002. The model values are overestimates and do not seem to have taken into account the observed EUV values.

INDEX TERMS: 7537 Solar Physics, Astrophysics, and Astronomy: Solar and stellar variability; 7507 Solar Physics, Astrophysics, and Astronomy: Chromosphere; 7509 Solar Physics, Astrophysics, and Astronomy: Corona; 7529 Solar Physics, Astrophysics, and Astronomy: Photosphere; 7534 Solar Physics, Astrophysics, and Astronomy: Radio emissions; **KEYWORDS:** solar indices, dissimilarities

Citation: Kane, R. P., Dissimilarity in the evolution of solar EUV and solar radio emission (2800 MHz) during 1999–2002, *J. Geophys. Res.*, 108(A12), 1455, doi:10.1029/2003JA009869, 2003.

1. Introduction

[2] Solar activity indices show variations in a wide range of time scales, from a few days to several years [Donnelly *et al.*, 1985, 1986; Hinteregger *et al.*, 1981]. Different indices originate in different parts of the solar atmosphere. Each solar region is also a source of radio emissions of certain frequencies, where higher frequencies escape from deeper regions. The magnitudes (percentage changes) differ considerably from one index to another [Lean, 1987]. In the work of Kane [2002a, 2002b], the behaviors of various indices near sunspot maxima and minima were compared for cycles 18–23. Based on the examination of 12-month moving averages, the general conclusion was that the evolutions of various solar indices around years of sunspot maxima and minima were not always similar to that of sunspots. Sometimes, lags and/or leads of a few months were seen.

[3] Since all solar indices attain maxima near sunspot maximum and minima near sunspot minimum, the long-term correlation between all indices was very high, particularly during rising and declining phases of a solar cycle. However, during the sunspot maximum years, dissimilarities of evolutions of different solar indices were often seen, even in 12-month moving averages. A particularly interesting feature of sunspot numbers was that during years around sunspot maxima, there was generally only one

prominent maximum, but in some cycles, there was a broad plateau. If the beginning and end of the plateau are termed as first and second maxima (separated by several months), the first maximum for sunspots was generally the higher one, and the valley in between was very shallow. Solar indices at or near the solar surface generally showed similar structures, with maxima matching with sunspot maxima within a month or two. Indices originating in the chromosphere and above showed two peaks in roughly the same months as sunspots, but for some indices, the second maximum was larger than the first maximum, and the valley between the two maxima was deeper, as compared with sunspot maxima.

[4] For several studies of Sun-Earth relationship, the variations of solar indices need to be established. For ionospheric F region, solar EUV is important. Observations of EUV have been very few (for technical reasons) and in the absence of EUV data, other solar indices are used as proxy. An index consistently used earlier as proxy for EUV is F10, the (observed) solar radio emission at 2800 MHz, 10.7 cm, mainly because the data for F10 are available regularly for a long time (since 1947). Sunspot data are also available, for a much longer time, but correlation studies indicate that in cases where EUV data are not available for comparison, ionospheric parameters are better correlated with F10 than with sunspots. Since 1981, all models, from the reference spectrum (SC#21REFW) of Hinteregger *et al.* [1981] to SOLAR2000 of Tobiska *et al.* [2000], have used F10 copiously. Since 1996, EUV measurements are available in a regular and reliable way [Judge *et al.*, 1998]. In the present communication, the

variations of F10 and EUV are compared during 1999–2002.

2. Data

[5] Sunspot and F10 data were obtained from the NOAA website (available at ftp://ftp.ngdc.noaa.gov/STP/SOLAR_DATA/), EUV data were obtained from the University of Southern California website [Judge *et al.*, 1998] (available at http://www.usc.edu/dept/space_science/semdata.htm), Lyman alpha data were obtained from the University of Colorado at Boulder ftp site [Woods *et al.*, 2000a] (available at ftp://laspftp.colorado.edu/pub/solstice/composite_lya.dat), Mg II data were obtained from the NOAA website (available at <http://www.sec.noaa.gov/ftpdir/sbuvs/NOAAMgII.dat>), and SOLAR2000 model data were obtained from Space Environment Technology's Space Weather Division website (available at <http://spacewx.com/>).

3. Plots of Observed Values

[6] Figure 1a shows the plots of the observed monthly values for F10 (2800 MHz flux), two EUV ranges 26–34 nm and 0.1–50 nm, Lyman alpha, Mg II and sunspot number Rz. There are considerable month-to-month fluctuations (mostly irregular), but one can see a first maximum near July 2000 and a second maximum near February 2002 (both marked by dots). To bring out the maxima more clearly, 12-month running means were calculated and are shown in Figure 1b. The maxima are marked by dots. The numbers in parentheses below the dots are the actual values at the maxima, while the numbers in rectangles above the dots are normalized values at the maxima, with values for the first maximum set as 100. For Rz, the second maximum is lower than the first maximum, by $\sim 4\%$. For other indices, the second maximum is higher than the first maximum, but there are quantitative differences. For 2800 MHz (10.7 cm) flux (F10), the second maximum is higher than the first maximum by 9.4%, but for the two EUV ranges, the second maximum is higher only by $\sim 3.5\%$. For Lyman alpha, the second maximum is higher by $\sim 8.9\%$, almost the same as for F10 (9.4%). For Mg II, the second maximum is higher only by $\sim 1\%$. (It may be remembered, however, that changes in Mg II are always smaller. For example, from the solar minimum in 1996 to the solar maximum in 2000–2002, F10 and EUV increased by a factor of ~ 2.80 , a change of 180%, Lyman alpha by a factor of ~ 1.6 , a change of $\sim 60\%$, while Mg II increased by a factor of 1.16 only, a change of 16%; see details in the work of Kane [2002c]). Thus the relative heights of the first and second maxima are similar for F10 and Lyman alpha (second maximum larger by $\sim 9\%$) but much smaller for EUV (second maximum larger only by $\sim 3.5\%$). For sunspot number Rz, there is a qualitative difference (second maximum smaller, by $\sim 4\%$). An examination of several other indices (not shown here), indicated that for radio frequencies up to ~ 1500 MHz escaping from upper corona, the first maximum was higher, while for higher frequencies escaping from middle and lower corona (including 2800 MHz), the second maximum was higher. Indices near the photosphere had mostly the first maximum higher. In the work of Kane [2002c], it was shown that amplitudes are generally low near the photosphere,

increase in the upper chromosphere, diminish in the transition region (temperature $\sim 100,000$ K) and then increase rapidly in the lower corona, diminishing further in the upper corona. This pattern is similar to that of the “emission measure distribution” reported by Warren *et al.* [1998, 2001] and should have an implication for the physical model of EUV flux generation in solar dynamo system.

4. Comparison With the Irradiance Model Estimates of SOLAR2000

[7] As inputs in the terrestrial atmosphere, one needs Irradiance models. These involve (1) reference spectra, i.e., the flux versus wavelength, and (2) formulae for their variations with solar cycle. Donnelly and Pope [1973] gave a reference spectrum for moderate solar activity (F 10 = 150). Using the AE-E data, Hinteregger *et al.* [1981] provided a reference spectrum (SC#21REFW) for low solar activity (cycle 21 solar minimum, F 10 = 68). These were followed by the models of Nusinov [1984], Tobiska and Barth [1990], Schmidke *et al.* [1992], Tobiska [1991], Richards *et al.* [1994], and Tobiska and Eparvier [1998]. All these empirical solar EUV models were derived from the AE-E data. However, some accurate EUV measurements from sounding rockets during solar cycle 22 (1992–1994) indicated that the irradiances based on the AE-E data could be underestimated by as much as a factor of 2 at some wavelengths [Woods *et al.*, 2000b; Bailey *et al.*, 2000; Solomon *et al.*, 2001]. Hence a big collaborative project was planned, which started in 1998 and has resulted in SOLAR2000 [Tobiska *et al.*, 2000], which is a new image- and full-disk proxy empirical solar irradiance model, with 1 nm resolution in the spectral range 1–1,000,000 nm for historical modeling and forecasting. For use of research workers who would like to have a quantitative measure of solar input in the terrestrial atmosphere, a new solar proxy (termed E10.7) has been generated as an output product of the SOLAR2000 model. It is the time-dependent solar EUV flux at the top of the Earth's atmosphere, integrated over the range 1–105 nm, and expressed in 2800 MHz 10.7 cm (F10) radio flux units.

[8] From the Space Environment Technology's Space Weather Division website (available at <http://SpaceWx.com>) (Vers. 1.24 SOLAR2000RG link), we accessed the spectra for 39 wavelength intervals 1.86–2.95 nm, 3.00–4.92 nm, ..., 100.1–105 nm for each day during 1996–2002, obtained the monthly mean spectra, combined these for broader wavelength intervals, and obtained their 12-month moving averages. The plots of the Irradiance model for the two EUV ranges 26–34 nm and 1–50 nm are shown in Figure 1c. Both EUV ranges 26–34 nm and 1–50 nm show the second maximum higher than the first maximum but by a large margin, $\sim 20\%$, almost 5 times the margin for observed EUV ($\sim 3.5\%$). The third plot is for E10.7, the SOLAR2000 model average over the range 1–105 nm. Here too, the margin is large, $\sim 20\%$, same as for SOLAR2000 EUV. Thus SOLAR 2000 model values show the second maximum larger than the first maximum by a factor of 2 or more as compared with the observed values, contrary to the assertion that the model values track observed EUV values within $\sim 5\%$ (Kent Tobiska, private communication, 2001). Workers using the SOLAR2000 model values need to keep this difference in mind. The model values can




Figure 1. Plots for 1999–2002 of (a) the monthly values of six solar indices, namely F10 (2800 MHz flux), the two observed EUV ranges 26–34 nm and 0.1–50 nm, and observed Lyman alpha and Mg II, (b) 12-month running means of the same six solar indices, and (c) 12-month running means of the SOLAR2000 model values of EUV ranges 26–34 nm and 1–50 nm, and the parameter E10.7 (integrated model solar EUV flux 1–105 nm). The maxima peaks are marked with dots. The numbers in parentheses are values at the peaks (relative to 1.0 at solar minimum of 1996), and the numbers in rectangles are normalized values, with values at the first peak set as 100.

be different from the observed values (in the present case, overestimates), by as much as 15–20%. This is not a random statistical uncertainty but a systematic difference.

5. Conclusions

[9] For solar EUV, the observed F10 (2800 MHz, 10.7 cm solar radio emission) has been used copiously as a proxy.

Even in the latest Solar Irradiance model SOLAR2000, F10 seems to be an important factor. In this communication, it is shown that during 1999–2002, F10 and sunspot number Rz showed two maxima, one near July 2000 and another near January 2002, with a minimum in between near February 2001. The second maximum was higher than the first maximum by $\sim 10\%$ for F10 and Lyman alpha, but for sunspot number Rz, the second maximum was lower than

the first maximum by $\sim 4\%$. The observed EUV (26–24 nm from SEM-SOHO) had the second maximum higher than the first maximum but only by $\sim 3.5\%$. The model values given in SOLAR2000 [Tobiska *et al.*, 2000] show patterns of 26–34 nm with the second maximum higher than the first maximum by as much as $\sim 20\%$. Thus there is a discrepancy of $\sim 15\%$ between observed EUV and SOLAR2000 model EUV during 2001–2002. The model values show ratios of second to first maximum (20%) much larger than even those of observed F10 and Lyman alpha ($\sim 9\%$). The model values are overestimates by $\sim 15\%$, at least in the present case. Workers using the model values as inputs for any investigation need to keep this possibility in mind.

[10] **Acknowledgments.** Thanks are due to the various groups who obtained these data and made them available in websites. SOLAR2000 Research Grade V1.24 historical irradiances are provided courtesy of W. Kent Tobiska and SpaceWx. These historical irradiances have been developed with funding from the NASA UARS, TIMED, SOHO, and SDO missions, as well as the JSDAP and LWS programs. Our work was partially supported by FNDCT, Brazil under contract FINEP-537/CT.

[11] Shadia Rifai Habbal thanks W. Kent Tobiska and two other referees for their assistance in evaluating this paper.

References

- Bailey, S. M., T. N. Woods, C. A. Barth, S. C. Solomon, L. R. Canfield, and R. Korde, Measurements of the solar soft X-ray irradiance from the Student Nitric Oxide Explorer: First analysis and underflight calibrations, *J. Geophys. Res.*, *105*, 27,179–27,193, 2000.
- Donnelly, R. F., and J. H. Pope, The 1–3000 Å solar flux for a moderate level of solar activity for use in modeling the ionosphere and upper atmosphere, *Tech. Rep. ERL 276-SEL 25*, Natl. Ocean. and Atmos. Admin., Boulder, Colo., 1973.
- Donnelly, R. F., J. W. Harvey, D. F. Sheath, and T. P. Repoff, Temporal characteristics of the solar UV flux and HE I line at 1083 nm, *J. Geophys. Res.*, *90*, 6267–6273, 1985.
- Donnelly, R. F., H. E. Hinteregger, and D. F. Sheath, Temporal variations of solar EUV, UV, and 10,830-Å radiations, *J. Geophys. Res.*, *91*, 5567–5578, 1986.
- Hinteregger, H. E., K. Fukui, and B. G. Gilson, Observational, reference, and model data on solar EUV, from measurements on AE-E, *Geophys. Res. Lett.*, *8*, 1147–1150, 1981.
- Judge, D. L., *et al.*, First solar EUV irradiances obtained from SOHO by the CELIAS/SEM, *Solar Phys.*, *177*, 161–173, 1998.
- Kane, R. P., Evolutions of various solar indices around sunspot maximum and sunspot minimum years, *Ann. Geophys.*, *20*, 735–741, 2002a.
- Kane, R. P., Correlations of solar indices with solar EUV fluxes, *Solar Phys.*, *207*, 17–40, 2002b.
- Kane, R. P., Short-term and long-term variability of solar emissions in recent years, *J. Geophys. Res.*, *107*(A10), 1298, doi:10.1029/2001JA000290, 2002c.
- Lean, J., Solar ultraviolet irradiance variations: A review, *J. Geophys. Res.*, *92*, 839–868, 1987.
- Nusinov, A. A., Dependence of the intensity of lines of short-wavelength solar radiation on the solar activity level, *Geomagn. Aeron.*, *24*, 439–444, 1984.
- Richards, P. G., J. A. Fennelly, and D. G. Torr, EUVAC—A solar EUV flux model for aeronomic calculations, *J. Geophys. Res.*, *99*, 8981–8992, 1994.
- Schmidtke, G., T. Woods, J. Worden, H. Doll, S. C. Solomon, and G. Rottman, Solar EUV irradiance from San Marco ASSI: A reference spectrum, *Geophys. Res. Lett.*, *19*, 2175–2178, 1992.
- Solomon, S. C., S. M. Bailey, and T. N. Woods, Effect of solar soft X-rays on the lower ionosphere, *Geophys. Res. Lett.*, *28*, 2149–2152, 2001.
- Tobiska, W. K., Revised solar extreme ultraviolet flux model, *J. Atmos. Terr. Phys.*, *53*, 1005–1018, 1991.
- Tobiska, W. K., and C. A. Barth, A solar EUV flux model, *J. Geophys. Res.*, *95*, 8243–8251, 1990.
- Tobiska, W. K., and F. Eparvier, EUV97, Improvements to EUV irradiance modeling in the soft X-rays and FUV, *Solar Phys.*, *177*, 147–159, 1998.
- Tobiska, W. K., T. Woods, F. Eparvier, R. Viereck, L. Flyod, D. Bouwer, G. Rottman, and O. R. White, The SOLAR2000 empirical solar irradiance model and forecast tool, *J. Atmos. Sol. Terr. Phys.*, *62*, 1233–1250, 2000.
- Warren, H. P., J. T. Mariska, and J. Lean, A new reference spectrum for the EUV irradiance of the quiet Sun, *J. Geophys. Res.*, *103*, 12,077–12,089, 1998.
- Warren, H. P., J. T. Mariska, and J. Lean, A new model of solar EUV irradiance variability: 1. Model formulation, *J. Geophys. Res.*, *106*, 15,745–15,757, 2001.
- Woods, T. N., W. K. Tobiska, G. J. Rottman, and J. R. Worden, Improved solar Lyman alpha irradiance modeling from 1947 through 1999 based on UARS observations, *J. Geophys. Res.*, *105*, 27,195–27,215, 2000a.
- Woods, T. N., G. J. Rottman, and S. C. Solomon, Solar extreme ultraviolet irradiance measurements from sounding rockets during solar cycle 22, *Phys. Chem. Earth (C)*, *25*, 397–399, 2000b.

R. P. Kane, Instituto Nacional de Pesquisas Espaciais, Caixa Postal 515, 12201-970 São Jose dos Campos, São Paulo, Brazil. (kane@laser.inpe.br)